

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/26293>

Please be advised that this information was generated on 2017-12-05 and may be subject to change.

# Total Synthesis of Brevetoxin B. 1. CDEFG Framework

K. C. Nicolaou,\* E. A. Theodorakis, F. P. J. T. Rutjes, J. Tiebes, M. Sato, E. Untersteller, and X.-Y. Xiao

Department of Chemistry, The Scripps Research Institute  
10666 North Torrey Pines Road, La Jolla, California 92037  
Department of Chemistry and Biochemistry  
University of California, San Diego  
9500 Gilman Drive, La Jolla, California 92093

Received November 1, 1994

With its imposing structure, brevetoxin B (1), produced by *Gymnodinium breve* Davis, stood as a formidable challenge to synthetic chemists since its discovery and structural elucidation in 1981.<sup>1</sup> Brevetoxin's beautifully arranged molecular assembly includes 11 *trans*-fused rings, each containing an oxygen atom, with each fusion consisting of a C—C bond separating two adjacent ring oxygens and with all adjacent substituents flanking the oxygens placed *syn* to each other except on ring K. Its unprecedented architecture, its association with the "red tide" catastrophes,<sup>2</sup> and its potent neurotoxicity and interference with the function of sodium channels attracted serious attention from chemists<sup>3</sup> and biologists<sup>4</sup> alike. We now wish to announce, in this and the following communication,<sup>5</sup> the total synthesis of brevetoxin B (1) in its naturally occurring form.

Figure 1 outlines the strategic bond disconnections and retrosynthetic analysis of 1. The adopted strategy benefited from convergency (oxocene disconnections) and synthetic technologies developed in these laboratories specifically for constructing oxocene<sup>6</sup> and tetrahydropyran<sup>7</sup> systems.




Figure 1. Strategic bond disconnections and retrosynthetic analysis of brevetoxin B (1).

The construction of the CDEFG framework 4 described herein began with the previously reported intermediate 7 (Scheme 1).<sup>8</sup> Swern oxidation of 7 followed by a Wittig reaction with the appropriate reagent furnished, in 99% overall yield, compound 9 via aldehyde 8. Hydrogenation of 9 and selective, acid-induced monodesilylation gave alcohol 11 via 10 in 97% overall yield. Oxidation of 11 in a sequential fashion using Swern and NaClO<sub>2</sub> conditions resulted in carboxylic acid 12 (97%), which upon desilylation with TBAF led to 13 (91%). Lactonization of hydroxy acid 13 by the Yamaguchi method<sup>9</sup> and enol triflate formation gave 15 via 14 in 84% overall yield. Generation of the higher order cuprate derived from the lithio derivative of iodide 17a<sup>10</sup> and 17b followed by coupling<sup>11</sup> with triflate 15 and partial acid-induced orthoester hydrolysis resulted in formation of 18 via 16 (84% yield over two steps, *ca.* 2.4:1 ratio at C\* in favor of the desired isomer, *vide infra*). Regio- and stereoselective hydroboration of 18 followed by oxidative workup and alkaline hydrolysis furnished hydroxy acid 19 in 73% overall yield. Finally, lactonization<sup>9</sup> of 19 and separation of the C\* epimers afforded pure lactone 6 (60% yield, plus 25% of its C\* methyl epimer), whose structure was determined by X-ray crystallographic analysis (see ORTEP drawing of a derivative<sup>10</sup> of 6, Figure 2).

The fusion of the remaining three rings onto the DEFG system 6 to afford the targeted polycyclic framework 4 proceeded as depicted in Scheme 2. Thus, conversion of lactone 6 to its enol triflate (97%) followed by Cr/Ni-mediated coupling<sup>12</sup> with

(1) Lin, Y.-Y.; Risk, M.; Ray, S. M.; Van Engen, D.; Clardy, J.; Golik, J.; James, J. C.; Nakanishi, K. *J. Am. Chem. Soc.* 1981, 103, 6773. Lee, M. S.; Repeta, D. J.; Nakanishi, K.; Zagorski, M. G. *J. Am. Chem. Soc.* 1986, 108, 7855.

(2) Anderson, D. M. *Sci. Am.*, 1994, 8, 62 and references cited therein.

(3) Nicolaou, K. C.; Tiebes, J.; Theodorakis, E. A.; Rutjes, F. P. J. T.; Koide, K.; Sato, M.; Untersteller, E. *J. Am. Chem. Soc.* 1994, 116, 9371. Shimizu, Y. *Pure Appl. Chem.* 1982, 54, 1973. Yasumoto, T.; Murata, M. *Chem. Rev.* 1993, 93, 1897. Nakanishi, K. *Toxicon* 1985, 23, 473. Scheuer, P. J. *Tetrahedron* 1994, 50, 3. Kadota, I.; Matsukawa, Y.; Yamamoto, Y. *J. Chem. Soc., Chem. Commun.* 1993, 1638. Palazon, J. M.; Soler, M. A.; Ramirez, M. A.; Martin, V. S. *Tetrahedron Lett.* 1993, 34, 5467. Yamamoto, Y.; Yamada, J.; Kadota, I. *Tetrahedron Lett.* 1991, 32, 7069. Feng, F.; Murai, A. *Chem. Lett.* 1992, 1587. Alvarez, E.; Diaz, M. T.; Perez, R.; Ravelo, J. L.; Regeiro, A.; Vera, J. A.; Zurita, D.; Martin, J. D. *J. Org. Chem.* 1994, 59, 2848. For other selected articles from these laboratories, see: Reddy, K. R.; Skokotas, G.; Nicolaou, K. C. *Gazz. Chim. Ital.* 1993, 123, 337. Nicolaou, K. C. *Aldrichimica Acta* 1993, 26 (3), 62. Nicolaou, K. C.; Reddy, K. R.; Skokotas, G.; Sato, F.; Xiao, X.-Y.; Hwang, C.-K. *J. Am. Chem. Soc.* 1993, 115, 3558.

(4) *International Symposium on Red Tides*; Okaichi, T., Anderson, D. M., Nemoto, T., Eds.; Elsevier: New York, 1989. *Toxic Dinoflagellates*; Anderson, D. M., White, A. W., Baden, D. G., Eds.; Elsevier: Amsterdam, 1985. *Marine Toxins: Origin, Structure and Molecular Pharmacology*; Sherwood, H.; Strichartz, G., Eds.; ACS Symposium Series 418; American Chemical Society: Washington, DC, 1990. Rein, K. S.; Baden, D. G.; Gawley, R. E. *J. Org. Chem.* 1994, 59, 2101. Rein, K. S.; Lynn, B.; Gawley, R. E.; Baden, D. G. *J. Org. Chem.* 1994, 59, 2107. Baden, D. G.; Mende, T. J.; Szmant, A. M.; Trainer, V. L.; Edwards, R. A.; Roszell, L. E. *Toxicon* 1988, 26, 97. Poli, M. A.; Mende, T. J.; Baden, D. G. *Mol. Pharmacol.* 1986, 30, 129. Catterall, W. A. *Annu. Rev. Biochem.* 1986, 55, 953. Trainer, V. L.; Thomsen, W. J.; Catterall, W. A.; Baden, D. G. *Mol. Pharmacol.* 1991, 40, 988.

(5) Nicolaou, K. C.; Rutjes, F. P. J. T.; Theodorakis, E. A.; Tiebes, J.; Sato, M.; Untersteller, E. *J. Am. Chem. Soc.* 1995, 117, 1173.

(6) Nicolaou, K. C.; McGarry, D. G.; Somers, P. K.; Kim, B. H.; Ogilvie, W. W.; Yiannikouros, G.; Prasad, C. V. C.; Veale, C. A.; Hark, R. R. *J. Am. Chem. Soc.* 1990, 112, 6263. Nicolaou, K. C.; Prasad, C. V. C.; Hwang, C.-K.; Duggan, M. E.; Veale, C. A. *J. Am. Chem. Soc.* 1989, 111, 5321. Nicolaou, K. C.; Veale, C. A.; Hwang, C.-K.; Hutchinson, J.; Prasad, C. V. C.; Ogilvie, W. W. *Angew. Chem., Int. Ed. Engl.* 1991, 30, 299.

(7) Nicolaou, K. C.; Prasad, C. V. C.; Somers, P. K.; Hwang, C.-K. *J. Am. Chem. Soc.* 1989, 111, 5330. Nicolaou, K. C.; Duggan, M. E.; Hwang, C.-K. *J. Am. Chem. Soc.* 1989, 111, 6666. Nicolaou, K. C.; Duggan, M. E.; Hwang, C.-K.; Somers, P. K. *J. Chem. Soc., Chem. Commun.* 1985, 1359.


(8) Nicolaou, K. C.; Nugiel, D. A.; Couladouros, E.; Hwang, C.-K. *Tetrahedron* 1990, 46, 4517.

(9) Inanaga, J.; Hirata, K.; Saeki, H.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* 1979, 52, 1989.


(10) For preparation of and selected data for this compound, see the supplementary material.

(11) Tsushima, K.; Araki, K.; Murai, A. *Chem. Lett.* 1989, 1313. Lipshutz, B. H.; Sengupta, S. *Org. React. (N.Y.)* 1992, 41, 135.




Scheme 1. Construction of DEFG Ring System 6<sup>a</sup>

<sup>a</sup> Reagents and conditions: (a) 2.0 equiv of (COCl)<sub>2</sub>, 3.0 equiv of DMSO, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, then 7.0 equiv of Et<sub>3</sub>N, 0.5 h, 100%; (b) 2.0 equiv of TBSO(CH<sub>2</sub>)<sub>3</sub>PPh<sub>3</sub><sup>+</sup>I<sup>-</sup>, 1.5 equiv of NaHMDS, THF, 0 °C, 10 min, then 8, 0.5 h, 99%; (c) H<sub>2</sub>, 0.1 equiv of Pd/C (10%), 0.1 equiv of Na<sub>2</sub>CO<sub>3</sub>, EtOAc, 25 °C, 12 h, 100%; (d) 1.0 equiv of CSA, CH<sub>2</sub>Cl<sub>2</sub>/MeOH (1:1), 0 °C 1 h, 97%; (e) 2.0 equiv of (COCl)<sub>2</sub>, 3.0 equiv of DMSO, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, then 7.0 equiv of Et<sub>3</sub>N, 0.5 h; 1.5 equiv of NaClO<sub>2</sub>, 2.0 equiv of NaH<sub>2</sub>PO<sub>4</sub>, 2.0 equiv of 2-methyl-2-butene, *t*-BuOH/H<sub>2</sub>O (2:1), 25 °C, 1 h, 97%; (f) 5.0 equiv of TBAF, THF, 65 °C, 8 h, 91%; (g) 1.05 equiv of 2,4,6-trichlorobenzoyl chloride, 1.5 equiv of Et<sub>3</sub>N, THF, 0 °C, 2 h, then added to 5.0 equiv of DMAP, benzene (*c* = 0.05 mM), 80 °C, 1 h, 90%; (h) 5.0 equiv of LiHMDS, 1.5 equiv of HMPA, THF, -78 °C, 2 h, then 1.5 equiv of Tf<sub>2</sub>NPh, -78 → 25 °C, 93%; (i) 6.0 equiv of 17a, 10.0 equiv of *t*-BuLi, Et<sub>2</sub>O, -120 → -78 °C, 0.5 h, then 5.0 equiv of 17b, -78 → 30 °C, 0.5 h, Et<sub>2</sub>O/THF/HMPA (1:1:1), then 15, -78 → 0 °C, 2 h, 84%; (j) 0.3 equiv of PPTS, DME/H<sub>2</sub>O (1:1), 25 °C, 100%; (k) 6.0 equiv of BH<sub>3</sub>·THF, 0 °C, then 25 equiv of 3 N NaOH, 50 equiv of 30% H<sub>2</sub>O<sub>2</sub>, 89%; (l) 2.0 equiv of LiOH, DME/H<sub>2</sub>O (1:1), 25 °C, 82%; (m) 1.05 equiv of 2,4,6-trichlorobenzoyl chloride, 1.5 equiv of Et<sub>3</sub>N, THF, 0 °C, 2 h, then added to 5.0 equiv of DMAP, benzene (*c* = 0.05 mM), 80 °C, 1 h, 60% of 6, plus 25% of its C\* epimer (after column chromatography).

Figure 2. ORTEP of the bis(*p*-bromobenzoyl) derivative of 6.

aldehyde 20<sup>10</sup> furnished alcohol 21 (66%, mixture of epimers), which was deoxygenated via xanthate 22 (89%) by the Barton method<sup>13</sup> to afford 23 (67%). Regio- and stereospecific hydration of 23 via hydroboration/oxidation gave alcohol 24 (82%), which was silylated, leading to 25 (96%). A series of reactions involving DIBAL-H-mediated ester cleavage (98%), Dess–Martin oxidation (85%), Horner–Emmons olefination (99%), and acid-induced selective desilylation (100%) afforded α,β-unsaturated ester 5 via 26, 27 and 28. Exposure of 5 to KH led to the formation of the CDEFG ring system 29 in 90%

Scheme 2. Construction of CDEFG Ring System 4<sup>a</sup>

<sup>a</sup> Reagents and conditions: (a) 5.0 equiv of LiHMDS, 1.5 equiv of HMPA, THF, -78 °C, 2 h, then 1.5 equiv of Tf<sub>2</sub>NPh, -78 → 25 °C, 97%; (b) 6.0 equiv of 20, 6.0 equiv of CrCl<sub>2</sub>, 0.02 equiv of NiCl<sub>2</sub>, DMF, 25 °C, ultrasound, 3 h, 66%; (c) 3.0 equiv of CS<sub>2</sub>, 50.0 equiv of KH (added over 5 h), Et<sub>2</sub>O, then 10.0 equiv of MeI, 25 °C, 89%; (d) 4.0 equiv of *n*-Bu<sub>3</sub>SnH, 0.1 equiv of AIBN, benzene, 80 °C, 67%; (e) 5.0 equiv of BH<sub>3</sub>·THF, -30 °C, then 25 equiv of 3 N NaOH, 50 equiv of 30% H<sub>2</sub>O<sub>2</sub>, 82%; (f) 2.0 equiv of TESOTf, 2.5 equiv of 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>, -70 °C, 1 h, 96%; (g) 2.5 equiv of DIBAL-H, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, 5 min, 98%; (h) 1.7 equiv of Dess–Martin periodinane, CH<sub>2</sub>Cl<sub>2</sub>, 25 °C, 2 h, 85%; (i) 2.0 equiv of KHMDS, 0.2 equiv of 18-crown-6, 5.0 equiv of (MeO)<sub>2</sub>P(O)CH<sub>2</sub>CO<sub>2</sub>Me, THF, 0 °C, 0.5 h then add 27, 3 h, 99%; (j) 1.0 equiv of CSA, CH<sub>2</sub>Cl<sub>2</sub>/MeOH (2:1), 25 °C, 1 h, 100%; (k) 2.0 equiv of KH, THF, 25 °C, 2 h, 90%; (l) 1.3 equiv of DIBAL-H, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, 2 min, then 3.0 equiv of MeOH, 97%; (m) 2.0 equiv of Ph<sub>3</sub>PCHCO<sub>2</sub>Et, CH<sub>2</sub>Cl<sub>2</sub>, 25 °C, 12 h, 98%; (n) 2.5 equiv of DIBAL-H, CH<sub>2</sub>Cl<sub>2</sub>, -78 °C, 2 h, 96%; (o) 0.2 equiv of Ti(O<sup>*i*</sup>Pr)<sub>4</sub>, 0.2 equiv of (+)-diethyl tartrate, 2.0 equiv of *t*-BuOOH (5 N in decane), CH<sub>2</sub>Cl<sub>2</sub>, -20 °C, 5 h, 99%; (p) 5.0 equiv of SO<sub>3</sub>·pyridine, 10 equiv of Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>/DMSO (4:1), 0 °C; (q) 1.2 equiv of NaHMDS, 1.5 equiv of CH<sub>3</sub>PPh<sub>3</sub><sup>+</sup>Br<sup>-</sup>, THF, 25 °C, 1 h, 80% (over two steps); (r) 1.5 equiv of TBAF, THF, 25 °C, 3 h, 100%.

yield via a stereoselective Michael-type reaction.<sup>14</sup> Extension of the ester side chain via DIBAL-H reduction and phosphorane condensation furnished, via aldehyde 30 (97%), the α,β-unsaturated ester 31 (98%), which was reduced to allylic alcohol 32 (96%). Sharpless asymmetric epoxidation<sup>15</sup> of 32 using (+)-DET as the chiral auxiliary gave the corresponding hydroxy epoxide (99% yield), which was further oxidized to the aldehyde and subjected to a Wittig reaction to afford terminal olefin 33 (80% over two steps), and thence hydroxy epoxide 4 upon TBAF-induced desilylation (100%).

The elaboration of 4 to the ABCDEFG framework 3, the coupling of the latter to the IJK system 2 and the completion of the total synthesis of brevetoxin B (1) are described in the following communication.<sup>5,16</sup>

**Acknowledgment.** See following communication.<sup>5</sup>

**Supplementary Material Available:** See following communication.<sup>5</sup>

JA943553G

(12) Takai, K.; Tagashira, M.; Kuroda, T.; Oshima, K.; Utimoto, K.; Nozaki, H. *J. Am. Chem. Soc.* 1986, 108, 6048. Kishi, Y. *Pure Appl. Chem.* 1989, 61, 313.

(13) Barton, D. H. R.; McCombie, S. W. *J. Chem. Soc., Perkin Trans. I* 1975, 1574.

(14) Nicolaou, K. C.; Hwang, C.-K.; Duggan, M. E. *J. Am. Chem. Soc.* 1989, 111, 6682.

(15) Katsuki, T.; Sharpless, K. B. *J. Am. Chem. Soc.* 1980, 102, 5976.

(16) All new compounds exhibited satisfactory spectral and exact mass data. Yields refer to spectroscopically and chromatographically homogeneous materials.